

WO03051826

Publication Title:

3-PHENYL-2-ARYLALKYLTHIOPROPIONIC ACID DERIVATIVES AS
SELECTIVE AGONISTS OF PPAR-ALPHA

Abstract:

Abstract of WO03051826

A compound of formula (I); wherein R₁ represents chloro, fluoro or hydroxy as well as optical isomers and racemates thereof as well as pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, to processes for preparing such compounds, to their utility in treating clinical conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance, to methods for their therapeutic use and to pharmaceutical compositions containing them.

A compound of formula (I); wherein R represents chloro, fluoro or hydroxy as well as optical isomers and racemates thereof as well as pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof, to processes for preparing such compounds, to their utility in treating clinical conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance, to methods for their therapeutic use and to pharmaceutical compositions containing them. Data supplied from the esp@cenet database - Worldwide

Courtesy of <http://v3.espacenet.com>

**3-PHENYL-2-ARYLALKYLTHIOPROPIONIC ACID DERIVATIVES
AS SELECTIVE ANTAGONISTS OF PPAR-ALPHA**

Field of the invention

5 The present invention relates to certain novel 3-phenyl-2-arylalkylthiopropionic acid derivatives, to processes for preparing such compounds, to their the utility in treating clinical conditions including lipid disorders (dyslipidemias) whether or not associated with insulin resistance and other manifestations of the metabolic syndrome, to methods for their therapeutic use and to pharmaceutical compositions containing them.

10


Background of the invention

The metabolic syndrome including type 2 diabetes mellitus, refers to a cluster of manifestations including insulin resistance with accompanying hyperinsulinaemia, possibly
15 type 2 diabetes mellitus, arterial hypertension, central (visceral) obesity, dyslipidaemia observed as deranged lipoprotein levels typically characterised by elevated VLDL (very low density lipoproteins), small dense LDL particles and reduced HDL (high density lipoprotein) concentrations and reduced fibrinolysis.

20 Recent epidemiological research has documented that individuals with insulin resistance run a greatly increased risk of cardiovascular morbidity and mortality, notably suffering from myocardial infarction and stroke. In type 2 diabetes mellitus atherosclerosis related conditions cause up to 80% of all deaths.

25 In clinical medicine there is awareness of the need to increase the insulin sensitivity in patients with the metabolic syndrome and thus to correct the dyslipidaemia which is considered to cause the accelerated progress of atherosclerosis. However, currently this is not a universally accepted diagnosis with well-defined pharmacotherapeutic indications.

30 The S-enantiomer of the compound of formula C below




2-ethoxy-3-[4-(2-{4-methanesulfonyloxyphenyl}ethoxy)phenyl]propanoic acid, is disclosed in PCT Publication Number WO99/62872. This compound is reported to be a modulator of peroxisome proliferator-activated receptors (PPAR, for a review of the PPARs see T. M. Willson et al, J Med Chem 2000, Vol 43, 527) and has combined PPAR α /PPAR γ agonist activity (Structure, 2001, Vol 9, 699, P. Cronet et al). This compound is effective in treating conditions associated with insulin resistance.

Surprisingly a series of compounds has now been found which are selective PPAR α modulators.

Description of the invention

The present invention provides a compound of formula I



wherein R¹ represents chloro, fluoro or hydroxy as well as optical isomers and racemates thereof as well as pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof.

It will be appreciated by those skilled in the art the compounds of formula I contain an optically active centre and therefore can exist as enantiomers which can be separated as described later. It is expected that most, if not all, of the activity of the compounds of formula I resides in one enantiomer: either the S or the R enantiomer or the (+) or the (–)

enantiomer. The enantiomers which are more active in the assays which are described later are preferred forms of the present invention. It will be understood that the present invention includes all mixtures of this active enantiomer with the other enantiomer, for example the racemic mixture, which is a useful intermediate for the active enantiomer.

5

The active enantiomers may be isolated by separation of racemate for example by fractional crystallization, resolution or HPLC on a chiral column (for example a Chiralpak™ AD 250x50 column). Alternatively the active enantiomers may be made by chiral synthesis from chiral starting materials under conditions which will not cause racemisation or epimerisation, or by derivatisation with a chiral reagent.

10

The term "prodrug" as used in this specification includes derivatives of the carboxylic acid group which are converted in a mammal, particularly a human, into the carboxylic acid group or a salt or conjugate thereof. The term "prodrug" also includes derivatives of the hydroxy substituent (when R¹ represents hydroxy) which are converted in a mammal, particularly a human, into the hydroxy group or a salt or conjugate thereof. It should be understood that, whilst not being bound by theory, it is believed that most of the activity associated with the prodrugs arises from the activity of the compound of formula I into which the prodrugs are converted. Prodrugs can be prepared by routine methodology well within the capabilities of someone skilled in the art. Various prodrugs of carboxy and hydroxy are known in the art. For examples of such prodrug derivatives, see:

15

- a) Design of Prodrugs, edited by H. Bundgaard, (Elsevier, 1985) and Methods in Enzymology. 42: 309-396, edited by K. Widder, *et al.* (Academic Press, 1985);
- b) A Textbook of Drug Design and Development, edited by Krogsgaard-Larsen and H. Bundgaard, Chapter 5 "Design and Application of Prodrugs", by H. Bundgaard p.113-191 (1991);
- c) H. Bundgaard, Advanced Drug Delivery Reviews, 8:1-38 (1992);
- d) H. Bundgaard, *et al.*, Journal of Pharmaceutical Sciences, 77:285 (1988); and
- e) N. Kakeya, *et al.*, Chem Pharm Bull, 32:692 (1984).

20

The above documents a to e are herein incorporated by reference.

30

In vivo cleavable esters are just one type of prodrug of the parent molecule. An *in vivo* hydrolysable (or cleavable) ester of a compound of the formula (I) that contains a carboxy

or a hydroxy group is, for example, a pharmaceutically acceptable ester which is hydrolysed in the human or animal body to produce the parent acid or alcohol. Suitable pharmaceutically acceptable esters for carboxy include C₁₋₆alkoxymethyl esters, for example, methoxymethyl; C₁₋₆alkanoyloxymethyl esters, for example, pivaloyloxymethyl; phthalidyl esters; C₃₋₈cycloalkoxycarbonyloxyC₁₋₆alkyl esters, for example, 1-cyclohexyl-carbonyloxyethyl; 1,3-dioxolen-2-onylmethyl esters, for example, 5-methyl-1,3-dioxolen-2-onylmethyl; and C₁₋₆alkoxycarbonyloxyethyl esters, for example, 1-methoxycarbonyloxyethyl; and may be formed at any carboxy group in the compounds of this invention. An *in vivo* hydrolysable (or cleavable) ester of a compound of the formula (I) that contains a hydroxy group includes inorganic esters such as phosphate esters (including phosphoramidic cyclic esters) and α -acyloxyalkyl ethers and related compounds which as a result of the *in vivo* hydrolysis of the ester breakdown to give the parent hydroxy group/s. Examples of α -acyloxyalkyl ethers include acetoxymethoxy and 2,2-dimethylpropionyloxy-methoxy. A selection of *in vivo* hydrolysable ester forming groups for hydroxy include alkanoyl, benzoyl, phenylacetyl and substituted benzoyl and phenylacetyl, alkoxycarbonyl (to give alkyl carbonate esters), dialkylcarbamoyl and *N*-(dialkylaminoethyl)-*N*-alkylcarbamoyl (to give carbamates), dialkylaminoacetyl and carboxyacetyl. Examples of substituents on benzoyl include morpholino and piperazino linked from a ring nitrogen atom via a methylene group to the 3- or 4- position of the benzoyl ring.

The compounds of formula I have activity as medicaments, in particular the compounds of formula I are selective agonists of PPAR α , that is, their EC₅₀ for PPAR α is at least ten times lower than their respective EC₅₀ for PPAR γ wherein the EC₅₀s are measured and calculated as described in the assays later in this document. The compounds of formula I are potent and selective.

Specific compounds of the invention are:

2-{{[2-(4-chlorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid; preferably either the (R)- enantiomer or the (S)- enantiomer;

- 2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid; preferably either the (R)- enantiomer or the (S)- enantiomer;
- 2-([2-(4-fluorophenyl)ethyl]thio)-3-[4-(2-{4- [(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid; preferably either the (R)- enantiomer or the (S)- enantiomer;
- (-)-2-([2-(4-chlorophenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid and
- (-)-2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;
- and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof.

It will be understood by those skilled in the art that where (-) occurs in the above list that the compound indicated has a negative rotation when measured using the conditions and concentration described in the experimental section. It should be understood that salts of the parent acid are included even if a particular salt exhibits a (+) rotation provided that the absolute configuration of the salt is the same as the configuration of the (-)-parent acid.

Specific enantiomers of the invention are selected from one or more of the following:

- (R)- 2-([2-(4-chlorophenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;
- (S)-2-([2-(4-chlorophenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;
- (R)-2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;
- (S)-2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;
- (R)-2-([2-(4-fluorophenyl)ethyl]thio)-3-[4-(2-{4- [(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;
- (S)- 2-([2-(4-fluorophenyl)ethyl]thio)-3-[4-(2-{4- [(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;

(-)-2-{[2-(4-chlorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid and

(-)-2-{[2-(4-hydroxyphenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;

- 5 and pharmaceutically acceptable salts, solvates and crystalline forms thereof.

In the present specification the expression "pharmaceutically acceptable salts" is intended to define but is not limited to base salts such as the alkali metal salts, alkaline earth metal salts, ammonium salts, salts with basic amino acids, and salts with organic amines.

10

It will also be understood that certain compounds of the present invention may exist in solvated, for example hydrated, as well as unsolvated forms. It is to be understood that the present invention encompasses all such solvated forms. Certain compounds of the present invention may exist as tautomers. It is to be understood that the present invention


- 15 encompasses all such tautomers.

Methods of preparation

The compounds of the invention may be prepared as outlined below. However, the invention is not limited to these methods, the compounds may also be prepared as described for structurally related compounds in the prior art. The reactions can be carried out according to standard procedures or as described in the experimental section.

20


Compounds of formula I may be prepared by reacting a compound of formula II



25


in which R^1 is as previously defined and R^2 represents a protecting group for carboxylic hydroxy group as described in the standard text "Protective Groups in Organic Synthesis", 2nd Edition (1991) by Greene and Wuts, with a de-protecting reagent. The protecting group may also be a resin, such as Wang resin or 2-chlorotrityl chloride resin. Protecting groups
 5 may be removed in accordance to techniques which are well known to those skilled in the art. One such protecting group is where R^2 represents C_{1-6} alkoxy group or an arylalkoxy group eg benzyloxy, such that COR^2 represents an ester. Such esters can be reacted with a de-protecting reagent e.g. a hydrolysing agent, for example lithium hydroxide in a mixture of THF and water, at a temperature in the range of 0-100°C to give compounds of
 10 formula I.

Compounds of formula II may be prepared by reacting a compound of formula III



15

in which R^2 is as previously defined and X is a leaving group for example halo e.g. chloro with a compound of formula IV



20 in which R^1 is as previously defined or a protected hydroxy group for example R^1 is benzyloxy.

Compounds of formula III and IV may be prepared by methods described in the Examples or by analogous methods known to those skilled in the art.

Compounds of formula II, III and IV are believed to be novel and are claimed herein as useful intermediates in the preparation of compounds of formula I.

- 5 The compounds of the invention may be isolated from their reaction mixtures using conventional techniques.

Persons skilled in the art will appreciate that, in order to obtain compounds of the invention in an alternative and in some occasions, more convenient manner, the individual process
10 steps mentioned hereinbefore may be performed in different order, and/or the individual reactions may be performed at different stage in the overall route (i.e. chemical transformations may be performed upon different intermediates to those associated hereinbefore with a particular reaction).

- 15 The expression "inert solvent" refers to a solvent which does not react with the starting materials, reagents, intermediates or products in a manner which adversely affects the yield of the desired product.

Pharmaceutical preparations

20

The compounds of the invention will normally be administered via the oral, parenteral, intravenous, intramuscular, subcutaneous or in other injectable ways, buccal, rectal, vaginal, transdermal and/or nasal route and/or via inhalation, in the form of pharmaceutical preparations comprising the active ingredient either as a free acid, or a pharmaceutical
25 acceptable organic or inorganic base addition salt, in a pharmaceutically acceptable dosage form. Depending upon the disorder and patient to be treated and the route of administration, the compositions may be administered at varying doses.

Suitable daily doses of the compounds of the invention in therapeutical treatment of humans are about 0.0001-100 mg/kg body weight, preferably 0.001-10 mg/kg body weight.

- 5 Oral formulations are preferred particularly tablets or capsules which may be formulated by methods known to those skilled in the art to provide doses of the active compound in the range of 0.5mg to 500mg for example 1 mg, 3 mg, 5 mg, 10 mg, 25mg, 50mg, 100mg and 250mg.

According to a further aspect of the invention there is thus provided a pharmaceutical
10 formulation including any of the compounds of the invention, or pharmaceutically acceptable derivatives thereof, in admixture with pharmaceutically acceptable adjuvants, diluents and/or carriers.

Pharmacological properties

15 The present compounds of formula (I) are useful for the prophylaxis and/or treatment of clinical conditions associated with inherent or induced reduced sensitivity to insulin (insulin resistance) and associated metabolic disorders (also known as metabolic syndrome). These clinical conditions will include, but will not be limited to, general
20 obesity, abdominal obesity, arterial hypertension, hyperinsulinaemia, hyperglycaemia, type 2 diabetes and the dyslipidaemia characteristically appearing with insulin resistance. This dyslipidaemia, also known as the atherogenic lipoprotein profile, is characterised by moderately elevated non-esterified fatty acids, elevated very low density lipoprotein (VLDL) triglyceride rich particles, high Apo B levels, low high density lipoprotein (HDL)
25 levels associated with low apoAI particle levels and high Apo B levels in the presence of small, dense, low density lipoproteins (LDL) particles, phenotype B.

The compounds of the present invention are expected to be useful in treating patients with combined or mixed hyperlipidemias or various degrees of hypertriglyceridemias and
30 postprandial dyslipidemia with or without other manifestations of the metabolic syndrome.

Treatment with the present compounds is expected to lower the cardiovascular morbidity and mortality associated with atherosclerosis due to their antidyslipidaemic as well as antiinflammatory properties. The cardiovascular disease conditions include macro-
5 angiopathies of various internal organs causing myocardial infarction, congestive heart failure, cerebrovascular disease and peripheral arterial insufficiency of the lower extremities. Because of their insulin sensitizing effect the compounds of formula I are also expected to prevent or delay the development of type 2 diabetes from the metabolic syndrome and diabetes of pregnancy. Therefore the development of long-term
10 complications associated with chronic hyperglycaemia in diabetes mellitus such as the micro-angiopathies causing renal disease, retinal damage and peripheral vascular disease of the lower limbs are expected to be delayed. Furthermore the compounds may be useful in treatment of various conditions outside the cardiovascular system whether or not associated with insulin resistance, like polycystic ovarian syndrome, obesity, cancer and
15 states of inflammatory disease including neurodegenerative disorders such as mild cognitive impairment, Alzheimer's disease, Parkinson's disease and multiple sclerosis.

The compounds of the present invention are expected to be useful in controlling glucose levels in patients suffering from type 2 diabetes.

20

The present invention provides a method of treating or preventing dyslipidemias, the insulin resistance syndrome and/or metabolic disorders (as defined above) comprising the administration of a compound of formula I to a mammal (particularly a human) in need thereof.

25

The present invention provides a method of treating or preventing type 2 diabetes comprising the administration of an effective amount of a compound of formula I to a mammal (particularly a human) in need thereof.

30 In a further aspect the present invention provides the use of a compound of formula I as a medicament.

In a further aspect the present invention provides the use of a compound of formula I in the manufacture of a medicament for the treatment of insulin resistance and/or metabolic disorders.

5 Combination Therapy

The compounds of the invention may be combined with other therapeutic agents that are useful in the treatment of disorders associated with the development and progress of atherosclerosis such as hypertension, hyperlipidaemias, dyslipidaemias, diabetes and
10 obesity. The compounds of the invention may be combined with another therapeutic agent that decreases the ratio of LDL:HDL or an agent that causes a decrease in circulating levels of LDL-cholesterol. In patients with diabetes mellitus the compounds of the invention may also be combined with therapeutic agents used to treat complications related to micro-angiopathies.

15 The compounds of the invention may be used alongside other therapies for the treatment of metabolic syndrome or type 2 diabetes and its associated complications, these include biguanide drugs, for example metformin, phenformin and buformin, insulin (synthetic insulin analogues, amylin) and oral antihyperglycemics (these are divided into prandial
20 glucose regulators and alpha-glucosidase inhibitors). An example of an alpha-glucosidase inhibitor is acarbose or voglibose or miglitol. An example of a prandial glucose regulator is repaglinide or nateglinide.

In another aspect of the invention, the compound of formula I, or a pharmaceutically
25 acceptable salt, solvate, solvate of such a salt or a prodrug thereof, may be administered in association with another PPAR modulating agent. PPAR modulating agents include but are not limited to a PPAR alpha and/or gamma agonist, or pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof. Suitable PPAR alpha and/or gamma
30 agonists, pharmaceutically acceptable salts, solvates, solvates of such salts or prodrugs thereof are well known in the art. These include the compounds described in WO 01/12187, WO 01/12612, WO 99/62870, WO 99/62872, WO 99/62871, WO 98/57941, WO 01/40170, J Med Chem, 1996, 39, 665, Expert Opinion on Therapeutic Patents, 10 (5),

623-634 (in particular the compounds described in the patent applications listed on page 634) and J Med Chem, 2000, 43, 527 which are all incorporated herein by reference.

Particularly a PPAR alpha and/or gamma agonist refers to NN622/Ragaglitazar, BMS 298585, WY-14643, clofibrate, fenofibrate, bezafibrate, gemfibrozil and ciprofibrate; GW 9578, ciglitazone, troglitazone, pioglitazone, rosiglitazone, eglitazone, proglitazone, BRL-49634, KRP-297, JTT-501, SB 213068, GW 1929, GW 7845, GW 0207, L-796449, L-165041 and GW 2433. Particularly a PPAR alpha and/or gamma agonist refers to (S)-2-ethoxy-3-[4-(2-(4-methanesulphonyloxyphenyl)ethoxy)-phenyl]propanoic acid and pharmaceutically acceptable salts thereof.

In addition the combination of the invention may be used in conjunction with a sulfonylurea for example: glimepiride, glibenclamide (glyburide), gliclazide, glipizide, gliquidone, chlorpropamide, tolbutamide, acetohexamide, glycopyramide, carbutamide, glibonuride, glisoxepid, glybuthiazole, glibuzole, glyhexamide, glymidine, glypinamide, phenbutamide, tolcyclamide and tolazamide. Preferably the sulfonylurea is glimepiride or glibenclamide (glyburide). More preferably the sulfonylurea is glimepiride. Therefore the present invention includes administration of a compound of the present invention in conjunction with one, two or more existing therapies described in this paragraph. The doses of the other existing therapies for the treatment of type 2 diabetes and its associated complications will be those known in the art and approved for use by regulatory bodies for example the FDA and may be found in the Orange Book published by the FDA. Alternatively smaller doses may be used as a result of the benefits derived from the combination.

The present invention also includes a compound of the present invention in combination with a cholesterol-lowering agent. The cholesterol-lowering agents referred to in this application include but are not limited to inhibitors of HMG-CoA reductase (3-hydroxy-3-methylglutaryl coenzyme A reductase). Suitably the HMG-CoA reductase inhibitor is a statin selected from the group consisting of atorvastatin, bervastatin, cerivastatin, dalvastatin, fluvastatin, itavastatin, lovastatin, mevastatin, nicostatin, nivastatin, pravastatin and simvastatin, or a pharmaceutically acceptable salt, especially sodium or calcium, or a solvate thereof, or a solvate of such a salt. A particular statin is atorvastatin, or a

pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof. A more particular statin is atorvastatin calcium salt. A particularly preferred statin is, however, a compound with the chemical name (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl(methylsulfonyl)-amino]-pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid, [also known as (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[N-methyl-N-(methylsulfonyl)-amino]pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid] or a pharmaceutically acceptable salt or solvate thereof, or a solvate of such a salt. The compound (E)-7-[4-(4-fluorophenyl)-6-isopropyl-2-[methyl-(methylsulfonyl)-amino]-pyrimidin-5-yl](3R,5S)-3,5-dihydroxyhept-6-enoic acid, and its calcium and sodium salts are disclosed in European Patent Application, Publication No. EP-A-0521471, and in Bioorganic and Medicinal Chemistry, (1997), 5(2), 437-444. This latter statin is now known under its generic name rosuvastatin.

In the present application, the term "cholesterol-lowering agent" also includes chemical modifications of the HMG-CoA reductase inhibitors, such as esters, prodrugs and metabolites, whether active or inactive.

The present invention also includes a compound of the present invention in combination with an inhibitor of the ileal bile acid transport system (IBAT inhibitor).

Suitable compounds possessing IBAT inhibitory activity have been described, see for instance the compounds described in WO 93/16055, WO 94/18183, WO 94/18184, WO 96/05188, WO 96/08484, WO 96/16051, WO 97/33882, WO 98/07449, WO 98/03818, WO 98/38182, WO 99/32478, WO 99/35135, WO 98/40375, WO 99/35153, WO 99/64409, WO 99/64410, WO 00/01687, WO 00/47568, WO 00/61568, WO 00/62810, WO 01/68906, DE 19825804, WO 00/38725, WO 00/38726, WO 00/38727, WO 00/38728, WO 00/38729, WO 01/68906, WO 01/66533, WO 02/32428, WO 02/50051, EP 864 582, EP489423, EP549967, EP573848, EP624593, EP624594, EP624595 and EP624596 and the contents of these patent applications are incorporated herein by reference.

Particular classes of IBAT inhibitors suitable for use in the present invention are benzothiepinines, and the compounds described in the claims, particularly claim 1, of WO 00/01687, WO 96/08484 and WO 97/33882 are incorporated herein by reference. Other suitable classes of IBAT inhibitors are the 1,2-benzothiazepines, 1,4-benzothiazepines and 1,5-benzothiazepines. A further suitable class of IBAT inhibitors is the 1,2,5-benzothiadiazepines.

One particular suitable compound possessing IBAT inhibitory activity is (3*R*,5*R*)-3-butyl-3-ethyl-1,1-dioxo-5-phenyl-2,3,4,5-tetrahydro-1,4-benzothiazepin-8-yl β -D-glucopyranosiduronic acid (EP 864 582). Other suitable IBAT inhibitors include one of:

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(carboxymethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(carboxymethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(2-sulphoethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(2-sulphoethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(5-carboxypentyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 5 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-(2-sulphoethyl)carbamoyl]-2-fluorobenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 10 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{(R)-1-[*N''*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]-2-hydroxyethyl}carbamoyl)benzyl]carbamoylmethoxy)-
- 15 2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-(ethoxy)(methyl)phosphoryl-methyl}carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-
- 20 tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{2-[(hydroxy)(methyl)phosphoryl]ethyl}carbamoyl)benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-methylthio-1-
- 25 carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{2-[(methyl)(ethyl)phosphoryl]ethyl}carbamoyl)-4-hydroxybenzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-(N'-\{2-[(methyl)(hydroxy)phosphoryl]ethyl\}carbamoyl)-4-hydroxybenzyl]carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,5-benzothiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[(R)-N'-(2-methylsulphinyl-1-carboxyethyl)carbamoyl]benzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,5-benzothiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methoxy-8- $\{N-[(R)-\alpha-[N'-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,5-benzothiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((R)-1-carboxy-2-methylthio-ethyl)carbamoyl]-4-hydroxybenzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((S)-1-carboxy-2-(R)-hydroxypropyl)carbamoyl]-4-hydroxybenzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((S)-1-carboxy-2-methylpropyl)carbamoyl]-4-hydroxybenzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((S)-1-carboxybutyl)carbamoyl]-4-hydroxybenzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((S)-1-carboxypropyl)carbamoyl]benzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((S)-1-carboxyethyl)carbamoyl]benzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-((S)-1-carboxy-2-(R)-hydroxypropyl)carbamoyl]benzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8- $\{N-[(R)-\alpha-[N-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl\}carbamoylmethoxy\}-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;$

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxyethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((R)-1-carboxy-2-methylthioethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-{(S)-1-[*N*-((S)-2-hydroxy-1-carboxyethyl)carbamoyl]propyl}carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-methylpropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxypropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-[*N*-((R/S)- α -{*N*-[1-(R)-2-(S)-1-hydroxy-1-(3,4-dihydroxyphenyl)prop-2-yl]carbamoyl}-4-hydroxybenzyl)carbamoylmethoxy]-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine; and

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to an additional further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or

carrier, with the simultaneous, sequential or separate administration one or more of the following agents selected from:

a CETP (cholesteryl ester transfer protein) inhibitor, for example those referenced and described in WO 00/38725 page 7 line 22 - page 10, line 17 which are incorporated herein
5 by reference;

a cholesterol absorption antagonist for example azetidinones such as SCH 58235 and those described in US 5,767,115 which are incorporated herein by reference;

a MTP (microsomal transfer protein) inhibitor for example those described in Science, 282, 751-54, 1998 which are incorporated herein by reference;

10 a nicotinic acid derivative, including slow release and combination products, for example, nicotinic acid (niacin), acipimox and nicheritol;

a phytosterol compound for example stanols;

probucol;

an anti-obesity compound for example orlistat (EP 129,748) and sibutramine (GB

15 2,184,122 and US 4,929,629);

an antihypertensive compound for example an angiotensin converting enzyme (ACE)

inhibitor, an angiotensin II receptor antagonist, an adrenergic blocker, an alpha

adrenergic blocker, a beta adrenergic blocker, a mixed alpha/beta adrenergic blocker,

an adrenergic stimulant, calcium channel blocker, an AT-1 blocker, a saluretic, a diuretic

20 or a vasodilator;

a CB1 antagonist or inverse agonist for example as described in WO01/70700 and EP 65635 ;

a Melanin concentrating hormone (MCH) antagonist;

a PDK inhibitor; or

25 modulators of nuclear receptors for example LXR, FXR, RXR, and RORalpha;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof,

optionally together with a pharmaceutically acceptable diluent or carrier to a warm-

blooded animal, such as man in need of such therapeutic treatment.

30 Particular ACE inhibitors or pharmaceutically acceptable salts, solvates, solvate of such salts or a prodrugs thereof, including active metabolites, which can be used in combination with a compound of formula I include but are not limited to, the following compounds:

alacepril, alatriopril, altiopril calcium, ancovenin, benazepril, benazepril hydrochloride, benazeprilat, benzoylcaptopril, captopril, captopril-cysteine, captopril-glutathione, ceranapril, ceranopril, ceronapril, cilazapril, cilazaprilat, delapril, delapril-diacid, enalapril, enalaprilat, enapril, epicaptopril, foroxymithine, fosfenopril, fosenopril, fosenopril sodium, 5 fosinopril, fosinopril sodium, fosinoprilat, fosinoprilic acid, glycopril, hemorphan-4, idrapril, imidapril, indolapril, indolaprilat, libenzapril, lisinopril, lyciumin A, lyciumin B, mixanpril, moexipril, moexiprilat, moveltipril, muracein A, muracein B, muracein C, pentopril, perindopril, perindoprilat, pivalopril, pivopril, quinapril, quinapril hydrochloride, quinaprilat, ramipril, ramiprilat, spirapril, spirapril hydrochloride, spiraprilat, spiropril, 10 spiropril hydrochloride, temocapril, temocapril hydrochloride, teprotide, trandolapril, trandolaprilat, utibapril, zabicipril, zabiciprilat, zofenopril and zofenoprilat. Preferred ACE inhibitors for use in the present invention are ramipril, ramiprilat, lisinopril, enalapril and enalaprilat. More preferred ACE inhibitors for uses in the present invention are ramipril and ramiprilat.

15

Preferred angiotensin II antagonists, pharmaceutically acceptable salts, solvates, solvate of such salts or a prodrugs thereof for use in combination with a compound of formula I include, but are not limited to, compounds: candesartan, candesartan cilexetil, losartan, valsartan, irbesartan, tasosartan, telmisartan and eprosartan. Particularly preferred 20 angiotensin II antagonists or pharmaceutically acceptable derivatives thereof for use in the present invention are candesartan and candesartan cilexetil.

Therefore in an additional feature of the invention, there is provided a method for for the treatment of type 2 diabetes and its associated complications in a warm-blooded animal, 25 such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of one the other compounds described in this combination section, or a pharmaceutically acceptable salt, solvate, solvate of such a salt 30 or a prodrug thereof.

Therefore in an additional feature of the invention, there is provided a method of treating hyperlipidemic conditions in a warm-blooded animal, such as man, in need of such treatment which comprises administering to said animal an effective amount of a compound of formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof in simultaneous, sequential or separate administration with an effective amount of one the other compounds described in this combination section or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the invention there is provided a pharmaceutical composition which comprises a compound of formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and one of the other compounds described in this combination section or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in association with a pharmaceutically acceptable diluent or carrier.

According to a further aspect of the present invention there is provided a kit comprising a compound of formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and one of the other compounds described in this combination section or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a first unit dosage form;
- b) one of the other compounds described in this combination section or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof; in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to a further aspect of the present invention there is provided a kit comprising:

- a) a compound of formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, together with a pharmaceutically acceptable diluent or carrier, in a first unit dosage form;
- b) one of the other compounds described in this combination section or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in a second unit dosage form; and
- c) container means for containing said first and second dosage forms.

According to another feature of the invention there is provided the use of a compound of the formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and one of the other compounds described in this combination section, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the the treatment of metabolic syndrome or type 2 diabetes and its associated complications in a warm-blooded animal, such as man.

According to another feature of the invention there is provided the use of a compound of the formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, and one of the other compounds described in this combination section, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, in the manufacture of a medicament for use in the treatment of hyperlipidaemic conditions in a warm-blooded animal, such as man.

According to a further aspect of the present invention there is provided a combination treatment comprising the administration of an effective amount of a compound of the formula I, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier, with the simultaneous, sequential or separate administration of an effective amount of one of the other compounds described in this combination section, or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof, optionally together with a pharmaceutically acceptable diluent or carrier to a warm-blooded animal, such as man in need of such therapeutic treatment.

Examples

¹H NMR and ¹³C NMR measurements were performed on a Varian Mercury 300 or Varian UNITY plus 400, 500 or 600 spectrometers, operating at ¹H frequencies of 300, 400, 500 and 600 MHz, respectively, and at ¹³C frequencies of 75, 100, 125 and 150 MHz, respectively. Measurements were made on the delta scale (δ).

Unless otherwise stated, chemical shifts are given in ppm with the solvent as internal standard.

Abbreviations

DMSO	dimethyl sulfoxide
EtOAc	ethyl acetate
DMF	<i>N,N</i> -dimethylformamide
THF	tetrahydrofuran
MeCN	acetonitrile
MeOH	methanol
TFA	trifluoroacetic acid
NH ₄ OAc	ammonium acetate
t	triplet
s	singlet
d	doublet
q	quartet
m	multiplet
bs	broad singlet

Example 12-{[2-(4-Chlorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid5 (i) 2-(4-Chlorophenyl)ethyl]ethanethioate

1-(2-Chloroethyl)-4-chlorobenzene (0.700 g, 4.00 mmol), thioacetic acid (0.320 g, 4.20 mmol) and triethylamine (0.425 g, 4.20 mmol) were dissolved in MeOH (8 ml). The solution was warmed with microwaves for 900 seconds at 150 °C. The mixture was used in the next step without purification.

10

(ii) 2-{[2-(4-Chlorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid

The reaction vessel was flushed with argon. To a part 0.8 ml (0.400 mmol) of the mixture from the reaction step above was added under stirring sodium methanethiolate (56.5 mg, 0.800 mmol) in MeOH (0.2 ml). After an hour methyl 2-chloro-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate (0.200 g, 0.484 mmol, prepared as described in Example 2) in acetonitrile (0.800 ml) was added. The mixture was stirred for 16 h under an argon atmosphere and then centrifuged under vacuum at 35 °C. 2 ml 0.5 M LiOH solution (THF/water 7:1) was added and the mixture was stirred vigorously for 20 h. After acidification with 12 M HCl (100 µl) the stirring was continued for another hour. The crude product was filtered through a TeflonTM filter and further purified using preparative HPLC (0.2 % TFA/MeCN, gradient). The pooled fractions were centrifuged under vacuum giving 18 mg of the desired product (yield 7%).

¹H-NMR (400 MHz, CDCl₃): 2.75-2.95 (m, 5H), 3.06 (t, 2H), 3.10 (s, 3H), 3.16 (dd, 1H), 3.45 (dd, 1H), 4.14 (d, 2H), 6.87 (d, 2H), 7.05 (d, 2H), 7.11-7.27 (m, 8H).

(iii) (-)-2-{[2-(4-chlorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid

30 The (-)-enantiomer of 2-{[2-(4-chlorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid was separated from the racemate by chiral

chromatography. A Chiralpak AD JF003 (250 x 20 mm i.d.) column was used and ethanol/formic acid 100/0.1% as mobile phase. The racemate (2 g) was dissolved in ethanol (20 mg/ml) and was injected onto the column. The first eluting peak was collected and UV-detected. The product (0.9 g) was obtained with an enantiomeric purity = 99.2%.

- 5 The optical rotation was found to be $[\alpha]_D^{20} = -29^\circ$ by dissolving the enantiomer in ethanol to give a concentration of 0.5 g/100ml. The optical rotation was measured at 20 °C using the sodium line at 589 nm.

¹H NMR (500 MHz, CD₃OD): 7.18-7.25 (6H, m), 7.15 (2H, d), 7.13 (2H, d), 4.16 (2H, t), 3.45 (1H, t), 3.14 (3H, s), 3.11 (1H, m), 3.04 (2H, t), 2.77-2.90 (5H m).

10

Example 2

2-([2-(4-Hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid

- 15 (i) Methyl 2-chloro-3-[4-(2-hydroxyethyl)phenyl]propanoate

2-(4-Aminophenyl)ethanol (11g, 81mmol) and 32ml conc HCl was dissolved in acetone and cooled to 0°C. Sodium nitrite (5.6g, 81mmol) in 20ml water was added dropwise. The temperature was kept under 0°C. After one hour, methyl acrylate (70g, 808mmol) and CuI
20 (1.6g, 8mmol) were added (<0°C). The reaction mixture was stirred at room temperature overnight.

The solvent was evaporated and water was added. The water phase was extracted three times with EtOAc, the organic phases were pooled and washed with water, dried (MgSO₄) and evaporated under reduced pressure. The crude product was purified by flash
25 chromatography using a 65:35 mixture of EtOAc and heptane as eluent. Further purification by preparative HPLC (using a gradient of CH₃CN/ 5%CH₃CN-waterphase containing 0.1M NH₄OAc as eluent) gave 9.7g product (yield 49%) as an oil.

¹H NMR (400MHz, CDCl₃): 2.84 (t, 3H), 3.15 (dd, 1H), 3.35 (dd, 1H), 3.75 (s, 3H), 3.84 (t, 3H), 4.43 (t, 1H), 7.17 (d, 4H)

30

(ii) Methyl 3-(4-{2-[4-(benzyloxy)phenoxy]ethyl}phenyl)-2-chloropropanoate

Triphenylphosphine (2.4g, 9mmol) was added to a solution of methyl 2-chloro-3-[4-(2-hydroxyethyl)phenyl]propanoate (2.1g, 8.5mmol) and 4-(benzyloxy)phenol (1.7g, 8mmol) in 20ml toluene under nitrogen atmosphere. The solution was warmed to 55°C and diisopropyl azodicarboxylate (1.8g, 9mmol) was added. The reaction mixture was stirred at 55°C overnight.

The mixture was allowed to cool and the solvent was evaporated under reduced pressure. The crude product was purified by flash chromatography using a 80:20 mixture of heptane and EtOAc as eluent to yield 2.28g of the desired product (yield 61%) as colourless crystals.

¹HNMR (400MHz, CDCl₃): 3.05 (t, 2H), 3.16 (dd, 1H), 3.36 (dd, 1H), 3.75 (s, 3H), 4.12 (t, 2H), 4.45 (t, 1H), 5.01 (s, 2H), 6.82 (m, 2H), 6.90 (m, 2H), 7.13-7.27 (m, 4H), 7.29-7.47 (m, 5H).

(iii) Methyl 2-chloro-3-{4-[2-(4-hydroxyphenoxy)ethyl]phenyl}propanoate

Methyl 3-(4-{2-[4-(benzyloxy)phenoxy]ethyl}phenyl)-2-chloropropanoate (1.0g, 2.4mmol) and dimethyl sulfide (0.9g, 14mmol) was dissolved in 60ml CH₂Cl₂. Boron trifluoride etherate (2.0g, 14mmol) was added dropwise to the stirred solution. The reaction mixture was stirred for two days at room temperature. Another equivalent (0.4g, 2.87mmol) boron trifluoride etherate was added and the stirring was continued overnight. Water was added. The phases were separated and the aqueous phase was extracted twice with CH₂Cl₂. The organic phases were pooled, washed (water, brine), dried (Na₂SO₄) and evaporated under reduced pressure. Further purification by preparative HPLC using a gradient of CH₃CN/ 5% CH₃CN-waterphase containing 0.1M NH₄OAc gave 0.55g of the desired product (yield 52%) as an oil.

¹HNMR (400MHz, CDCl₃): 3.04 (t, 2H), 3.16 (dd, 1H), 3.35 (dd, 1H), 3.75 (s, 3H), 4.10 (t, 2H), 4.40 (t, 1H), 6.75 (m, 4H), 7.12-7.29 (m, 4H).

(iv) Methyl 2-chloro-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate

Methyl 2-chloro-3-[4-[2-(4-hydroxyphenoxy)ethyl]phenyl]propanoate (334mg, 1.0mmol) and triethylamine (303mg, 3.0mmol) was dissolved in 20ml dichlormethane and cooled to
 5 -20°C under nitrogen atmosphere. Methanesulfonyl chloride (114mg, 1.0mmol) was added dropwise. The mixture was allowed to reach room temperature. After 2 hours dichlormethane was added, the mixture was washed (water, brine), dried (Na₂SO₄) and evaporated under reduced pressure to yield 394mg pure product (yield 96%).

¹HNMR (400MHz, CDCl₃): 3.02-3.11 (m, 5H), 3.15 (dd, 1H), 3.35 (dd, 1H), 3.74 (s, 3H),
 10 4.14 (t, 2H), 4.44 (t, 1H), 5.29 (s, 2H), 6.88 (d, 2H), 7.14-7.25 (m, 6H).

(v) Methyl 2-({2-[4-(benzyloxy)phenyl]ethyl}thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate

2-[4-(Benzyloxy)phenyl]ethanethiol (334mg, 1.4mmol), methyl 2-chloro-3-[4-(2-{4-
 15 [(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate (394mg, 0.95mmol) and potassium carbonate (189mg, 1.4mmol) were dissolved in 14ml dry DMF and stirred under nitrogen atmosphere at room temperature overnight.

The solvent was evaporated under reduced pressure and the residue was dissolved in toluene. The organic phase was washed (water, brine), dried (MgSO₄) and evaporated.
 20 Further purification by preparative HPLC using a gradient of CH₃CN/5% CH₃CN-waterphase containing 0.1M NH₄OAc gave 477mg of the desired product (yield 75%).
¹HNMR (400MHz, CDCl₃): 2.76-2.89 (m, 4H), 2.95 (dd, 1H), 3.09 (m, 5H), 3.20 (dd, 1H), 3.53 (m, 1H), 3.70 (s, 3H), 4.15 (t, 2H), 5.06 (s, 2H), 6.91 (m, 4H), 7.07-7.24 (m, 8H), 7.31-7.48 (m, 5H).

25

(vi) Methyl 2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate

To a solution of methyl 2-({2-[4-(benzyloxy)phenyl]ethyl}thio)-3-[4-(2-{4-
 30 [(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate (477mg, 0.8mmol) and 15ml dichlormethane, dimethyl sulfide (239mg, 3.8mol) and boron trifluoride etherate (545mg,

3.8mmol) were added. After 18 hours of stirring water was added to the reaction. The phases were separated and the aqueous phase was extracted twice with dichlormethane. The organic phases were pooled, dried (MgSO₄) and evaporated under reduced pressure. 274mg of the desired product (yield 67%) was obtained as an oil.

5 ¹HNMR (400MHz, CDCl₃): 2.70-2.85 (m, 4H), 2.91 (dd, 1H), 3.05 (t, 2H), 3.10 (s, 3H), 3.17 (dd, 1H), 3.49 (m, 1H), 3.68 (s, 3H), 4.13 (t, 2H), 6.72 (d, 2H), 6.87 (d, 2H), 6.99 (d, 2H), 7.10-7.22 (m, 6H)

(vii) 2-([2-(4-Hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid

10

Methyl 2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoate (105mg, 0.2mmol) was dissolved in 6.5ml of a 7:1 mixture of THF and water and cooled on an ice-bath. Lithium hydroxide (9.4mg, 0.4mmol) was added. Water was added to the reaction mixture after 24 hours of stirring at room temperature. The THF was evaporated under reduced pressure and the residue was acidified with 1M hydrochloric acid. The water phase was extracted with EtOAc (x3), the organic phases were pooled, washed (water, brine), dried (MgSO₄) and evaporated. The crude product was purified using preparative HPLC (eluent: CH₃CN / 5% CH₃CN-waterphase containing 0.1M NH₄OAc) to give 74mg of the desired product (yield 97%) as an oil.

20

¹HNMR (400MHz, CDCl₃): 2.68-2.95 (m, 5H), 3.05 (t, 2H), 3.10 (s, 3H), 3.17 (dd, 1H), 3.47 (m, 1H), 4.12 (t, 2H), 6.70 (d, 2H), 6.86 (d, 2H), 6.97 (d, 2H), 7.12-7.21 (m, 6H).

25 ¹³CNMR (100MHz, CDCl₃): 33.8, 35.1, 35.5, 37.2, 37.3, 48.1, 69.3, 115.6, 115.8, 123.3, 129.3, 129.4, 129.9, 132.3, 136.2, 136.9, 142.8, 154.4, 158.0, 177.2.

(viii) (-)-2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid

30

The racemate of 2-([2-(4-hydroxyphenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}ethyl)phenyl]propanoic acid was separated into its enantiomers using chiral

chromatography. A Chiralpak AD JDB01+ AS003 (336 x 100 mm i.d.) and ethanol/formic acid 100/0.01% was used as mobile phase. The racemate (9 g) was dissolved in ethanol and injected onto the column. The first eluting peak was collected and UV-detected. The product (4.1 g) was obtained with an enantiomeric purity >99%. The optical rotation was found to be $[\alpha]_D^{20} = -33^\circ$ by dissolving the enantiomer in methanol to give a concentration of 0.64 g/100ml. The optical rotation was measured at 20 °C using the sodium line at 589 nm.

¹H NMR (500 MHz, CD₃OD): 7.17-7.22 (6H, m), 6.99 (2H, d), 6.94 (2H, d), 6.69 (2H, d), 4.17 (2H, t), 3.46 (1H, t), 3.16 (3H, s), 3.13 (1H, dd), 3.05 (2H, t), 2.69-2.88 (5H, m).

Example 3

2-{[2-(4-Fluorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}-ethyl)phenyl]propanoic acid

(i) [2-(4-fluorophenyl)ethyl]ethanethioate

To a solution of DMF (60ml) and thioacetic acid (2.1g, 28mmol), cooled in an ice-bath, was added cesium carbonate (9.1g, 28mmol). The reaction mixture immediately turned orange. 1-(2-Chloroethyl)-4-fluorobenzene (4g, 25mmol) was added, the ice-bath was removed and the reaction was stirred overnight. The solvent was evaporated and the residue was dissolved in EtOAc. The organic phase was washed (water), dried (NaSO₄) and evaporated under reduced pressure to yield 5g (99%) of the desired product as an oil. ¹H NMR (400MHz, CDCl₃): 2.33 (s, 3H), 2.83 (t, 2H), 3.09 (t, 2H), 6.98 (t, 2H), 7.17 (m, 2H).

¹³C NMR (100MHz, CDCl₃): 30.8, 30.9, 35.3, 115.4, 115.6, 130.2, 130.3, 135.9, 160.7, 195.8.

(ii) Methyl 2-{[2-(4-fluorophenyl)ethyl]thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]phenoxy}-ethyl)phenyl]propanoate

[2-(4-Fluorophenyl)ethyl]ethanethioate (96mg, 0.48mmol) was dissolved in 1ml methanol under nitrogen atmosphere. To this slurry was added sodium methanethiolate (33.9mg,

0.48mmol). After 30 minutes of stirring methyl 2-chloro-3-[4-(2-{4-[(methylsulfonyl)-oxy]phenoxy}ethyl)phenyl]propanoate (200mg, 0.48mmol) dissolved in 2ml CH₃CN was added. The resulting mixture was stirred overnight.

The solvent was evaporated and the residue was dissolved in EtOAc. The organic phase
5 was washed (water), dried (MgSO₄) and evaporated under reduced pressure.

Further purification by preparative HPLC (using a gradient of CH₃CN / 5%CH₃CN-waterphase containing 0.1M NH₄OAc) gave 82mg of the desired product (yield 32%).

¹HNMR (500MHz, CDCl₃): 2.82-3.00 (m, 5H), 3.10 (t, 2H), 3.14 (s, 3H), 3.22 (dd, 1H),
3.53 (m, 1H), 3.72 (s, 3H), 4.18 (t, 2H), 6.92 (d, 2H), 7.0 (t, 2H), 7.15 (m, 4H), 7.23 (m,
10 4H).

(iii) 2-([2-(4-Fluorophenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid

15 Methyl 2-([2-(4-fluorophenyl)ethyl]thio)-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoate (77mg, 0.14mmol) was dissolved in 2.5ml of a 4:1 mixture of THF and water and cooled on an ice-bath. Lithium hydroxide (6.9mg, 0.29mmol) was added. Water was added after 2 days of stirring at room temperature. The THF was evaporated under reduced pressure. The aqueous phase was acidified with 1M
20 HCl and extracted with EtOAc three times. The organic phases were pooled, washed (water, brine), dried (MgSO₄) and evaporated. The crude product was purified using preparative HPLC (eluent: CH₃CN / 5% CH₃CN-waterphase containing 0.1M NH₄OAc). 24mg of the desired product (yield 30%) was obtained as an oil.

¹HNMR (500MHz, CDCl₃): 2.82-3.00 (m, 5H), 3.10 (t, 2H), 3.14 (s, 3H), 3.21 (dd, 1H),
25 3.52 (m, 1H), 4.18 (t, 2H), 6.90 (d, 2H), 6.99 (t, 2H), 7.13 (m, 2H), 7.16-7.26 (m, 6H).

Biological activity

Formulations

Compounds were dissolved in DMSO to obtain 16 mM stock solutions. Before assays, stock solutions were further diluted in DMSO and culture media.

5

GENERAL CHEMICALS AND REAGENTS

Luciferase assay reagent was purchased from Packard, USA. Restriction Enzymes were from Boehringer and Vent polymerase from New England Biolabs.

10 CELL LINES AND CELL CULTURE CONDITIONS

U2-OS, (Osteogenic sarcoma, Human) was purchased from ATCC, USA. Cells were expanded and refrozen in batches from passage number six. Cells were cultured in Dulbecco's modified Eagle medium (DMEM) with 25 mM glucose, 2 mM glutamine or 4 mM L-alanyl-L-glutamine, 10% fetal calf serum, at 5% CO₂. Phosphate buffered saline (PBS) without addition of calcium or magnesium was used. All cell culture reagents were from Gibco (USA) and 96-well cell culture plates were purchased from Wallach.

15

PLASMID CONSTRUCTS FOR HETEROLOGOUS EXPRESSION

Standard recombinant DNA techniques were carried out as described by Ausubel (7). The Luciferase reporter vector, pGL5UAS (clone consists of five copies of the GAL4 DNA binding sequence, 5'-CGACGGAGTACTGTCCTCCGAGCT-3', cloned into the SacI/XhoI sites of pGL3-Promoter (Promega). The SacI/XhoI fragment carrying the UAS sites was constructed using annealed overlapping oligonucleotides.

20

Expression vectors used are based upon pSG5 (Stratagene). All vectors contain an EcoRI/NheI fragment encoding the DNA binding domain of GAL4 (encoding amino acid positions 1-145 of database accession number P04386) followed by an in-frame fusion to a fragment encoding the nuclear localisation sequence from T antigen of Polyoma Virus. The nuclear localisation sequence was constructed using annealed overlapping

25

oligonucleotides creating *NheI*/*KpnI* sticky ends

(5'-CTAGCGCTCCTAGAAGAAACGCAAGGTTGGTAC-3'). The ligand binding

domains from human and mouse PPAR α and human and mouse PPAR γ were PCR amplified as *KpnI*/*BamHI* fragments and cloned in frame to the GAL4 DNA binding

5 domain and the nuclear localisation sequence. The sequence of all plasmid constructs used were confirmed by sequencing.

The following expression vectors were used for transient transfections:

vector	encoded PPAR subtype	sequence reference ¹
pSGGALhPPa	human PPAR α	S74349, nt 625-1530
pSGGALmPPa	murine PPAR α	X57638, nt 668-1573
pSGGALhPPg	human PPAR γ	U63415, nt 613-1518
pSGGALmPPg	murine PPAR γ	U09138, nt 652-1577

10 ¹ refers to nucleotide positions of data base entry used to express the ligand binding domain.

TRANSIENT TRANSFECTIONS

Frozen stocks of cells from passage number six were thawed and expanded to passage

15 number eight before transfections. Confluent cells were trypsinised, washed and pelleted by centrifugation at 270xg for 2 minutes. The cell pellet was resuspended in cold PBS to a

cell concentration of about 18×10^6 cells/ml. After addition of DNA, the cell suspension

was incubated on ice for approximately 5 minutes before electroporation at 230 V, 960 μ F in Biorad's Gene PulserTM in 0.5 ml batches. A total of 50 μ g DNA was added to each

20 batch of 0.5 ml cells, including 2.5 μ g expression vector, 25 μ g reporter vector and 22.5 μ g unspecific DNA (pBluescript, Stratagene).

After electroporation, cells were diluted to a concentration of 320'000 cells/ml in DMEM without phenol red, and approximately 25'000 cells/well were seeded in 96-well plates. In order to allow cells to recover, seeded plates were incubated at 37°C for 3-4 hours before addition of test compounds. In assays for PPAR α , the cell medium was supplemented with
5 resin-charcoal stripped fetal calf serum (FCS) in order to avoid background activation by fatty acid components of the FCS. The resin-charcoal stripped FCS was produced as follows; for 500 ml of heat-inactivated FCS, 10 g charcoal and 25 g Bio-Rad Analytical Grade Anion Exchange Resin 200-400 mesh were added, and the solution was kept on a magnetic stirrer at room temperature over night. The following day, the FCS was
10 centrifuged and the stripping procedure was repeated for 4-6 hours. After the second treatment, the FCS was centrifuged and filter sterilised in order to remove remnants of charcoal and resin.

ASSAY PROCEDURE

15 Stock solutions of compounds in DMSO were diluted in appropriate concentration ranges in master plates. From master plates, compounds were diluted in culture media to obtain test compound solutions for final doses.

After adjustment of the amount of cell medium to 75 μ l in each well, 50 μ l test compound
20 solution was added. Transiently transfected cells were exposed to compounds for about 24 hours before the luciferase detection assay was performed. For luciferase assays, 100 μ l of assay reagent was added manually to each well and plates were left for approximately 20 minutes in order to allow lysis of the cells. After lysis, luciferase activity was measured in a 1420 Multiwell counter, Victor, from Wallach.

25

Reference compounds

The TZD pioglitazone was used as reference substance for activation of both human and murine PPAR γ . 5,8,11,14-Eicosatetrayonic acid (ETYA) was used as reference substance for human PPAR α .

Calculations and analysis

For calculation of EC_{50} values, a concentration-effect curve was established. Values used were derived from the average of two or three independent measurements (after subtraction
5 of the background average value) and were expressed as the percentage of the maximal activation obtained by the reference compound. Values were plotted against the logarithm of the test compound concentration. EC_{50} values were estimated by linear intercalation between the data points and calculating the concentration required to achieve 50% of the maximal activation obtained by the reference compound.

10


The compounds of formula I have an EC_{50} of less than $5\mu\text{mol/l}$ for $PPAR\alpha$ and particular compounds have an EC_{50} of less than $1.0\mu\text{mol/l}$. Additionally in particular compounds the ratio of the EC_{50} ($PPAR\gamma$) : EC_{50} ($PPAR\alpha$) is greater than 25:1. It is believed that this ratio is important with respect to the pharmacological activity of the compounds and to
15 their therapeutic profile.

In addition the compounds of the present invention exhibit improved DMPK (Drug Metabolism and Pharmacokinetic) properties for example they exhibit improved metabolic stability *in vitro*. The compounds also have a promising toxicological profile.

20

CLAIMS

1. A compound of formula I



wherein R^1 represents chloro, fluoro or hydroxy as well as optical isomers and racemates thereof as well as pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof.

2. A compound selected from

2-{{2-(4-chlorophenyl)ethyl}thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;

2-{{2-(4-hydroxyphenyl)ethyl}thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid; and


2-{{2-(4-fluorophenyl)ethyl}thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid;

and pharmaceutically acceptable salts, prodrugs, solvates and crystalline forms thereof.

3. The compound (-)-2-{{2-(4-chlorophenyl)ethyl}thio}-3-[4-(2-{4-[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid.


4. The compound (-)-2-{[2-(4-hydroxyphenyl)ethyl]thio}-3-[4-(2-{4-
[(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid.
- 5 5. The (R)- enantiomer or the (S)- enantiomer of 2-{[2-(4-fluorophenyl)ethyl]thio}-3-[4-
(2-{4- [(methylsulfonyl)oxy]-phenoxy}ethyl)phenyl]propanoic acid.
6. A pharmaceutical formulation comprising a compound according to any one of claims
1 to 5 in admixture with pharmaceutically acceptable adjuvants, diluents and/or carriers.
- 10 7. A pharmaceutical formulation comprising a compound according to any one of claims
1-5 in admixture with pharmaceutically acceptable adjuvants, diluents and/or carriers.
8. A method of treating or preventing lipid disorders (dyslipidemia) whether or not
15 associated with insulin resistance comprising the administration of a compound according
to any one of claims 1-5 to a mammal in need thereof.
9. The use of a compound according to any one of claims 1-5 in the manufacture of a
medicament for the treatment of lipid disorders (dyslipidemia) whether or not associated
20 with insulin resistance.
10. A method of treating or preventing type 2 diabetes comprising the administration of an
effective amount of a compound of formula I according to any one of claims 1-5 to a
mammal in need thereof.
- 25 11. A process to prepare compounds of formula I reacting a compound of formula II

36




in which R^1 is as previously defined and represents R^2 represents a protecting group for carboxylic hydroxy group with a de-protecting reagent.

5 12. A compound of formula II



wherein R^1 is as previously described and R^2 represents a protecting group for carboxylic hydroxy.

10 13. A compound of formula III



in which R^2 is as previously defined and X is a leaving group .

15

14. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 5 combined with another therapeutic agent that is useful in the treatment of

disorders associated with the development and progress of atherosclerosis such as hypertension, hyperlipidaemias, dyslipidaemias, diabetes and obesity.

15 15. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 5 combined with another PPAR modulating agent.

16. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 5 combined with a cholesterol-lowering agent.

10 17. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 5 combined with a HMG-CoA reductase inhibitor.

18. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 5 combined with atorvastatin or a pharmaceutically acceptable salt, solvate,
15 crystalline form or prodrug thereof.

19. A pharmaceutical composition comprising a compound as claimed in any one of claims 1 to 5 combined with rosuvastatin or a pharmaceutically acceptable salt thereof.

20. A pharmaceutical composition comprising a compound as claimed in any one of
20 claims 1 to 5 combined with an IBAT inhibitor.

21. A pharmaceutical composition according to claim 20 wherein the IBAT inhibitor is selected from one of:

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(carboxymethyl)
25 carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(carboxymethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(2-sulphoethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
30

- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)-1'-phenyl-1'-[*N'*-(2-sulphoethyl)carbamoyl]methyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(5-carboxypentyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-2-fluorobenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-{*N*-[(R)- α -(*N'*-{(R)-1-[*N''*-(R)-(2-hydroxy-1-carboxyethyl)carbamoyl]-2-hydroxyethyl}carbamoyl)benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;
- 1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(carboxymethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-{ α -[*N'*-

((ethoxy)(methyl)phosphoryl-methyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3-butyl-3-ethyl-5-phenyl-7-methylthio-8-(*N*-[(*R*)- α -(*N'*-{2-

5 [(hydroxy)(methyl)phosphoryl]ethyl}carbamoyl]benzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N'*-(2-methylthio-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

10 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-[(R)- α -(*N'*-{2-[(methyl)(ethyl)phosphoryl]ethyl}carbamoyl)-4-hydroxybenzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-[(R)- α -(*N'*-{2-[(methyl)(hydroxy)phosphoryl]ethyl}carbamoyl)-4-hydroxybenzyl]carbamoylmethoxy)-2,3,4,5-tetrahydro-

15 1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[(R)-*N'*-(2-methylsulphinyl-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

20 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methoxy-8-(*N*-{(R)- α -[*N'*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,5-benzothiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((R)-1-carboxy-2-methylthio-ethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

25 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-(R)-hydroxypropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-methylpropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxybutyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxypropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxyethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-(R)-hydroxypropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-(2-sulphoethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxyethyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((R)-1-carboxy-2-methylthioethyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-{(S)-1-[*N*-((S)-2-hydroxy-1-carboxyethyl)carbamoyl]propyl}carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxy-2-methylpropyl)carbamoyl]benzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-((S)-1-carboxypropyl)carbamoyl]-4-hydroxybenzyl}carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;
- 1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-[*N*-((R/S)- α -{*N*-[1-(R)-2-(S)-1-hydroxy-1-(3,4-dihydroxyphenyl)prop-2-yl]carbamoyl}-4-hydroxybenzyl)carbamoylmethoxy]-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]-4-hydroxybenzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine; and

1,1-Dioxo-3,3-dibutyl-5-phenyl-7-methylthio-8-(*N*-{(R)- α -[*N*-(2-(S)-3-(R)-4-(R)-5-(R)-2,3,4,5,6-pentahydroxyhexyl)carbamoyl]benzyl} carbamoylmethoxy)-2,3,4,5-tetrahydro-1,2,5-benzothiadiazepine;

or a pharmaceutically acceptable salt, solvate, solvate of such a salt or a prodrug thereof.

INTERNATIONAL SEARCH REPORT

Intern Application No
PCT/EP 02/05743

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C323/56 A61K31/192

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, WPI Data, PAJ, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 62872 A (ASTRA) 9 December 1999 (1999-12-09) cited in the application page 4; claims 1,5-11 -----	1-13
A	WO 99 62871 A (ASTRA) 9 December 1999 (1999-12-09) examples 1,43,44 -----	1-13

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *G* document member of the same patent family

Date of the actual completion of the international search

7 May 2003

Date of mailing of the international search report

22/05/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

English, R.

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-21 (in part)

The scope of claims 1-21, in as far as the expressions "prodrug", "protecting group" and "leaving group" are concerned, is so unclear (Article 6 PCT) that a meaningful International Search is not possible with regard to these expressions. Nevertheless the search was complete for compounds of claims 12 and 13 in which the protecting group R2 is a methoxy group and for compounds of claim 13 in which the leaving group is a chlorine atom.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO Guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

Intel ional application No.
PCT/GB 02/05743

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although claims 8,10 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☒ Claims Nos.: 1-21 (in part)
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Internat Application No
PCT/GB 02/05743

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9962872	A	09-12-1999	AU 752261 B2	12-09-2002
			AU 4667199 A	20-12-1999
			AU 752262 B2	12-09-2002
			AU 4667299 A	20-12-1999
			BR 9910921 A	06-03-2001
			BR 9910928 A	13-02-2001
			CA 2333938 A1	09-12-1999
			CA 2334374 A1	09-12-1999
			CN 1311772 T	05-09-2001
			CN 1312795 T	12-09-2001
			EE 200000720 A	15-04-2002
			EE 200000725 A	17-06-2002
			EP 1084103 A1	21-03-2001
			EP 1084102 A1	21-03-2001
			HR 20000782 A1	30-06-2001
			HU 0103226 A2	28-01-2002
			HU 0103376 A2	29-05-2002
			JP 2002516899 T	11-06-2002
			JP 2002516900 T	11-06-2002
			NO 20006115 A	07-02-2001
			NO 20006116 A	02-02-2001
			PL 344681 A1	19-11-2001
			PL 345205 A1	03-12-2001
			WO 9962872 A1	09-12-1999
			WO 9962871 A1	09-12-1999
			SK 17682000 A3	06-08-2001
			SK 17692000 A3	10-05-2001
			TR 200003581 T2	21-06-2001
			TR 200003583 T2	21-05-2001
			TW 446696 B	21-07-2001
			US 2003027859 A1	06-02-2003
			US 6258850 B1	10-07-2001
			US 2001034371 A1	25-10-2001
WO 9962871	A	09-12-1999	AU 4667099 A	20-12-1999
			AU 752262 B2	12-09-2002
			AU 4667299 A	20-12-1999
			BR 9910913 A	06-03-2001
			BR 9910921 A	06-03-2001
			CA 2334107 A1	09-12-1999
			CA 2334374 A1	09-12-1999
			CN 1312795 T	12-09-2001
			CN 1311769 T	05-09-2001
			EE 200000717 A	15-08-2001
			EE 200000725 A	17-06-2002
			EP 1084101 A1	21-03-2001
			EP 1084102 A1	21-03-2001
			HU 0103226 A2	28-01-2002
			JP 2002516898 T	11-06-2002
			JP 2002516899 T	11-06-2002
			NO 20006114 A	02-02-2001
			NO 20006116 A	02-02-2001
			PL 344681 A1	19-11-2001
			PL 344682 A1	19-11-2001
			WO 9962870 A1	09-12-1999
			WO 9962871 A1	09-12-1999
			SK 17672000 A3	06-08-2001
			SK 17692000 A3	10-05-2001

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/JP 02/05743

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9962871 A		TR 200003543 T2	20-04-2001
		TR 200003583 T2	21-05-2001
		TW 446694 B	21-07-2001
		ZA 200006771 A	20-05-2002
		AU 752261 B2	12-09-2002
		AU 4667199 A	20-12-1999
		BR 9910928 A	13-02-2001
		CA 2333938 A1	09-12-1999
		CN 1311772 T	05-09-2001
		EE 200000720 A	15-04-2002
		EP 1084103 A1	21-03-2001
		HR 20000782 A1	30-06-2001
		HU 0103376 A2	29-05-2002
		JP 2002516900 T	11-06-2002
		NO 20006115 A	07-02-2001
		PL 345205 A1	03-12-2001
		WO 9962872 A1	09-12-1999
		SK 17682000 A3	06-08-2001
		TR 200003581 T2	21-06-2001
		TW 446696 B	21-07-2001